

Start Saving Today with the NUCYNTA[®] ER/NUCYNTA[®] Savings Card



Pay as little as \$35* for each prescription

Pharmacy Locator Service 888-884-2655

*Offer valid for commercially insured patients only. Maximum savings limit applies; patient out-of-pocket expense may vary. I am in agreement with the Program Terms and Conditions received with this Co-pay Card. Use of this Co-pay Card confirms that I meet all offer eligibility criteria.

Please see Important Safety Information throughout this brochure. Please see full Prescribing Information for Nucynta[®] ER or Nucynta[®], including Boxed Warning and Medication Guide, in pocket of brochure or at Nucynta.com

BIN: 600426
PCN: 54
GROUP: EC14703002
ID: 49155122449

PROGRAM TERMS, CONDITIONS, AND ELIGIBILITY CRITERIA

Offer is valid for commercially insured patients only. Most eligible commercially insured patients will pay as little as \$35 (maximum benefit amount of \$100) for NUCYNTA ER or NUCYNTA tablets. Any remaining out-of-pocket expense will be the patient's responsibility. This offer is valid in the United States. Some restrictions apply. Offer not valid for prescriptions reimbursed under Medicaid, a Medicare drug benefit plan, Tricare or other federal or state health programs (such as medical assistance programs). Cash Discount Cards and other non-insurance plans are not valid as primary under this offer. If the patients for drug benefits under each any such program, the patient cannot use this offer. By using this offer, the patient certifies that he or she will comply with any terms of his or her health insurance contract requiring notification to his or her payer of the existence and/or value of this offer. This offer is not valid for cash paying patients. Offer not valid for patients under 18 years of age. **Program not valid in Massachusetts or in any other state or jurisdiction where prohibited or restricted by law.**

Patient Instructions: To redeem this offer, you must have a valid prescription for NUCYNTA ER or NUCYNTA and follow the dosage instructions provided by your healthcare provider with a valid Prescriber ID# (eg, NPI or DEA). This offer may not be redeemed for cash. Patients with questions about the NUCYNTA ER or NUCYNTA savings offer should call 1-844-807-0348.

Pharmacist: When you apply this offer, you are certifying that you have not submitted a claim for reimbursement under any federal, state, or other governmental programs for this prescription. Participation in this program must comply with all applicable laws and regulations as a pharmacy provider.

Pharmacist instructions for a patient with an Eligible Third Party Payer: Submit the claim to the primary Third Party Payer first, then submit the balance due to **Change Healthcare** as a Secondary Payer COB [coordination of benefits] with patient responsibility amount and a valid Other Coverage Code (**eg, 8**) NUCYNTA ER or NUCYNTA tablets. Reimbursement will be received from **Change Healthcare**. Valid Other Coverage Code required. For any questions regarding **Change Healthcare** online processing, please call the Help Desk at 1-800-433-4893.

Program expires 06/30/2020.

This offer is not transferable and is limited to one offer per person. Not valid if reproduced. Void where prohibited by law. Program managed by ConnectiveRx on behalf of Collegium Pharmaceutical. The parties reserve the right to rescind, revoke or amend this offer without notice at any time.

Please see full [Prescribing information](#), including [Boxed Warning](#), and [Medication Guide](#), at Nucynta.com.

Important Safety Information for NUCYNTA® ER (tapentadol)

NUCYNTA® ER (tapentadol) is:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily, around-the-clock, long-term treatment with an opioid when other pain treatments, such as non-opioid pain medicines or immediate-release opioid medicines, do not treat your pain well enough or you cannot tolerate them
- Also used to manage pain from damaged nerves (neuropathic pain) that happens with diabetes and is severe enough to require daily, around-the-clock, long-term treatment with an opioid when other pain treatments, such as non-opioid pain medicines, do not treat your pain well enough or you cannot tolerate them
- A long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed, you are at risk for opioid addiction, abuse, and misuse that can lead to death
- Not used to treat pain that is not around-the-clock pain

IMPORTANT SAFETY INFORMATION ABOUT NUCYNTA ER

WARNING: ADDICTION, ABUSE, AND MISUSE; RISK EVALUATION AND MITIGATION STRATEGY (REMS); LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; INTERACTION WITH ALCOHOL and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

Addiction, Abuse, and Misuse

NUCYNTA ER exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing NUCYNTA ER, and monitor all patients regularly for the development of these behaviors and conditions.

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a REMS for these products. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to

- complete a REMS-compliant education program,
- counsel patients and/or their caregivers, with every prescription, on safe use, serious risks, storage, and disposal of these products,
- emphasize to patients and their caregivers the importance of reading the Medication Guide every time it is provided by their pharmacist, and
- consider other tools to improve patient, household, and community safety.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of NUCYNTA ER. Monitor for respiratory depression, especially during initiation of NUCYNTA ER or following a dose increase. Instruct patients to swallow NUCYNTA ER tablets whole; crushing, chewing, or dissolving NUCYNTA ER tablets can cause rapid release and absorption of a potentially fatal dose of tapentadol.

Accidental Ingestion

Accidental ingestion of even one dose of NUCYNTA ER, especially by children, can result in a fatal overdose of tapentadol.

Neonatal Opioid Withdrawal Syndrome

Prolonged use of NUCYNTA ER during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available.

Interaction With Alcohol

Instruct patients not to consume alcoholic beverages or use prescription or non-prescription products that contain alcohol while taking NUCYNTA ER. The co-ingestion of alcohol with NUCYNTA ER may result in increased plasma tapentadol levels and a potentially fatal overdose of tapentadol.

Risks From Concomitant Use With Benzodiazepines or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death.

- Reserve concomitant prescribing of NUCYNTA ER and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.
- Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.

Important information about NUCYNTA ER:

- Get emergency help right away if you take too much NUCYNTA ER (overdose). When you first start taking NUCYNTA ER, when your dose is changed, or if you take too much (overdose), serious or life threatening breathing problems that can lead to death may occur
- Taking NUCYNTA ER with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death
- Never give anyone else your NUCYNTA ER tablets. They could die from taking it. Selling or giving away NUCYNTA ER Tablets is against the law
- Store NUCYNTA ER Tablets securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home

Do not take NUCYNTA ER if you have:

- severe asthma, trouble breathing, or other lung problems
- a bowel blockage or have narrowing of the stomach or intestines

Important Safety Information for NUCYNTA® ER (tapentadol) continued

Before taking NUCYNTA ER, tell your healthcare provider if you have a history of:

- head injury, seizures
- problems urinating
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.
- liver, kidney, thyroid problems
- pancreas or gallbladder problems

Tell your healthcare provider if you are:

- **pregnant or planning to become pregnant.** Prolonged use of NUCYNTA ER during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated
- **breastfeeding.** Not recommended during treatment with NUCYNTA ER. It may harm your baby
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking NUCYNTA ER with certain other medicines can cause serious side effects

When taking NUCYNTA ER:

- Do not change your dose. Take NUCYNTA ER exactly as prescribed by your healthcare provider. Use the lowest effective dose for the shortest time needed
- Take your prescribed dose every 12 hours, at the same time every day. Do not take more than your prescribed dose in 24 hours. If you miss a dose, take your next dose at your usual time
- Swallow NUCYNTA ER whole. Do not cut, break, chew, crush, dissolve, snort, or inject NUCYNTA ER because this may cause you to overdose and die
- **Call your healthcare provider if the dose you are taking does not control your pain**
- **Do not stop taking NUCYNTA ER without talking to your healthcare provider**
- Dispose of expired, unwanted, or unused NUCYNTA ER tablets by promptly flushing down the toilet, if a drug take-back option is not readily available. Visit www.fda.gov/drugdisposal for additional information on disposal of unused medicines

While taking NUCYNTA ER, DO NOT:

- Drive or operate heavy machinery until you know how NUCYNTA ER affects you. NUCYNTA ER can make you sleepy, dizzy, or lightheaded
- Drink alcohol, or use prescription or over-the-counter medicines containing alcohol. Using products containing alcohol during treatment with NUCYNTA ER may cause you to overdose and die

The possible side effects of NUCYNTA ER are:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe

Get emergency medical help if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, lightheadedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion
- agitation, hallucinations, coma, feeling overheated, or heavy sweating

These are not all the possible side effects of NUCYNTA ER. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information, go to dailymed.nlm.nih.gov**

Please see full [Prescribing information](#), including [Boxed Warning](#), and [Medication Guide](#), at Nucynta.com, or speak to your healthcare provider if you have questions about NUCYNTA ER.

Important Safety Information for NUCYNTA® (tapentadol)

NUCYNTA tablets are:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage short term (acute) pain in adults when other pain treatments such as non-opioid pain medicines do not treat your pain well enough or you cannot tolerate them
- An opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed, you are at risk for opioid addiction, abuse, and misuse that can lead to death

IMPORTANT SAFETY INFORMATION ABOUT NUCYNTA TABLETS

WARNING: ADDICTION, ABUSE, AND MISUSE; RISK EVALUATION AND MITIGATION STRATEGY (REMS); LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

Addiction, Abuse, and Misuse

NUCYNTA tablets expose patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing NUCYNTA tablets, and monitor all patients regularly for the development of these behaviors and conditions.

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a REMS for these products. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to

- complete a REMS-compliant education program,
- counsel patients and/or their caregivers, with every prescription, on safe use, serious risks, storage, and disposal of these products,
- emphasize to patients and their caregivers the importance of reading the Medication Guide every time it is provided by their pharmacist, and
- consider other tools to improve patient, household, and community safety.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of NUCYNTA tablets. Monitor for respiratory depression, especially during initiation of NUCYNTA tablets or following a dose increase.

Accidental Ingestion

Accidental ingestion of even one dose of NUCYNTA tablets, especially by children, can result in a fatal overdose of tapentadol.

Neonatal Opioid Withdrawal Syndrome

Prolonged use of NUCYNTA tablets during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available.

Risks From Concomitant Use With Benzodiazepines or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death.

- Reserve concomitant prescribing of NUCYNTA tablets and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.
- Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.

Important information about NUCYNTA tablets:

- **Get emergency help right away if you take too much NUCYNTA tablets (overdose).** When you first start taking NUCYNTA tablets, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur
- Taking NUCYNTA tablets with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death
- Never give anyone else your NUCYNTA tablets. They could die from taking it. Selling or giving away NUCYNTA tablets is against the law
- Store NUCYNTA tablets securely, out of sight and reach of children, and in a location not accessible by others including visitors to the home

Do not take NUCYNTA tablets if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.

Before taking NUCYNTA tablets, tell your healthcare provider if you have a history of:

- head injury, seizures
- problems urinating
- abuse of street or prescription drugs, alcohol addiction, or mental health problems
- liver, kidney, thyroid problems
- pancreas or gallbladder problems

Tell your healthcare provider if you are:

- **pregnant or planning to become pregnant.** Prolonged use of NUCYNTA tablets during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated
- **breastfeeding.** NUCYNTA tablets pass into breast milk and may harm your baby
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking NUCYNTA tablets with certain other medicines can cause serious side effects that could lead to death

Important Safety Information for NUCYNTA® (tapentadol) continued

When taking NUCYNTA tablets:

- Do not change your dose. Take NUCYNTA tablets exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed
- Take your prescribed dose every 4-6 hours, at the same time every day. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.
- Call your healthcare provider if the dose you are taking does not control your pain
- If you have been taking NUCYNTA tablets regularly, do not stop taking NUCYNTA tablets without talking to your healthcare provider
- Dispose of expired, unwanted, or unused NUCYNTA tablets by promptly flushing down the toilet, if a drug take-back option is not readily available. Visit www.fda.gov/drugdisposal for additional information on disposal of unused medicines

While taking NUCYNTA tablets, DO NOT:

- Drive or operate heavy machinery until you know how NUCYNTA tablets affect you. NUCYNTA tablets can make you sleepy, dizzy, or lightheaded
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with NUCYNTA tablets may cause you to overdose and die

The possible side effects of NUCYNTA tablets:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe

Get emergency medical help if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, lightheadedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion

These are not all the possible side effects of NUCYNTA tablets. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information, go to dailymed.nlm.nih.gov**

Please see full [Prescribing information](#), including [Boxed Warning](#), and [Medication Guide](#), at Nucynta.com, or speak to your healthcare provider if you have questions about NUCYNTA tablets.